

ERRATA

Corrections to published RIFE reports

	Page, Section	Comment															
RIFE-1 1995	38, Section 16.2	Last but one sentence, replace 1994 with 1995.															
	39, Section 16.4	1 st sentence, 2 nd paragraph, replace 1994 with 1995.															
	45, Table 1	Replace ²⁴¹ Am Sellafield (sea pipelines) limit of 1.3 TBq with 0.3 TBq. Replace ⁶⁰ Co Harwell (pipeline) percentage of 1.5 with 6.9.															
	74, Table 16 99, Table 33(a)	The following activity in soil data were reported as being Bq kg ⁻¹ (dry) whilst they should have been reported as Bq kg ⁻¹ (wet). All data are averages unless stated.															
		<table border="1"> <thead> <tr> <th>Site/location</th> <th>²¹⁰Po</th> <th>²³⁸Pu</th> <th>²³⁹⁺²⁴⁰Pu</th> </tr> </thead> <tbody> <tr> <td>Sellafield (Table 16)</td> <td>64</td> <td></td> <td></td> </tr> <tr> <td>Aldermaston (Table 33(a))</td> <td></td> <td>0.0091</td> <td>0.36</td> </tr> <tr> <td>max</td> <td></td> <td>0.016</td> <td>0.56</td> </tr> </tbody> </table>	Site/location	²¹⁰ Po	²³⁸ Pu	²³⁹⁺²⁴⁰ Pu	Sellafield (Table 16)	64			Aldermaston (Table 33(a))		0.0091	0.36	max		0.016
Site/location	²¹⁰ Po	²³⁸ Pu	²³⁹⁺²⁴⁰ Pu														
Sellafield (Table 16)	64																
Aldermaston (Table 33(a))		0.0091	0.36														
max		0.016	0.56														
	99, Table 33(a)	The concentration of ¹³⁷ Cs in clay at Outfall (Pangbourne) was 12±0.15 Bq kg ⁻¹ (dry)															
	133, Appendix 3	The average consumption rates of nuts and offal by 10 year old children were 1.5 kg y ⁻¹ . The consumption of whelks at Sellafield by group E (Whitehaven commercial) was 11 kg y ⁻¹ .															
	138, Appendix 6	The values of t _f and t _s were 0. The transfer factors for beef offal (²⁴¹ Pu) and lamb (²⁴¹ Pu) were 2 10 ⁻² and 4 10 ⁻⁴ respectively.															
RIFE-2 1996	32, Section 8.1	lines 8-11. Replace with “In 1996 no fragments of spent fuel were found on the public beach at Dounreay. Thirteen small fragments were found with caesium-137 activities in the range 10 ⁵ -10 ⁸ Bq (these activities were measured by the operator). They were all found on the Dounreay foreshore which although a public area is largely inaccessible. A”															
	58, Table 2	Replace ³⁵ S Oldbury limit of 0.8 TBq with 0.75 TBq. Replace ⁴¹ Ar Trawsfynydd limit of 350 TBq with 3500 TBq.															

85, Table 16
87, Table 18
91, Table 20(a)
95, Table 21
119, Table 41

The following activity in soil data were reported as being Bq kg⁻¹ (dry) whilst they should have been reported as Bq kg⁻¹ (wet). All data are averages unless stated.

Site/location	²³⁴ U	²³⁵ U	²³⁸ U
Drigg (Table 16)	8.3	0.28	7.4
Ravenglass (Table 18)	16	0.56	15
Springfields (Table 20(a))	49	2.3	45
Capenhurst (Table 21)	9.8	0.36	10
Derby (Table 41)	44	1.7	43

Table 47

This was omitted in error. The data are attached.

Table 47. Radioactivity in plants near landfill sites, 1996

Sampling location	Material	No of samples	Mean radioactivity concentration (dry)*, Bq kg ⁻¹							
			³ H	¹⁴ C	⁹⁰ Sr	¹²⁵ I	¹³⁴ Cs	¹³⁷ Cs	²³⁸ Pu	²³⁹⁺²⁴⁰ Pu
Beddingham Lewes, East Sussex	Grass	4	<40 ±18	130 ±28	1.8 ±0.1	<0.19	<0.61	<0.54 ±0.30	<0.00099 ±0.00037	0.0067 ±0.0012
Cilgwyn Quarry, Gwynedd	"	4	<30	360 ±55	3.0 ±0.2	<0.63	<0.69	<5.2 ±0.9	<0.0095	0.018 ±0.005
Lyndown, Devon	"	4	<28	150 ±30	2.4 ±0.2	<1.3 ±0.2	<0.60	<0.62 ±0.17	<0.0010	<0.0024 ±0.0009
Witton, Cheshire	"	4	<38	130 ±33	0.76 ±0.12	<1.1 ±0.3	<0.59	<0.63	<0.0013	0.0021 ±0.0016

* Results are available for other artificial nuclides detectable by gamma spectrometry
All such results are less than the limit of detection

RIFE-3
1997

19, Table 1.1

Replace beta, tritium and ⁶⁰Co Devonport (sewer) discharges with 1.97 10⁻⁶, 2.22 10⁻⁶, 5.60 10⁻⁷ TBq respectively.
Replace alpha and beta limit and percentage Greenwich with 4.44 10⁻³ TBq and <1 respectively.

21, Table 1.2

Replace tritium Winfrith limit with 5 TBq.

38, Section 3.6.5

1st paragraph. Reference to factor of 0.85 millisievert per milligray should be ICRP (1996b).

70, Table 4.10
72, Table 4.12
81, Table 4.16
121, Table 9.1

The following activity in soil data were reported as being Bq kg⁻¹ (dry) whilst they should have been reported as Bq kg⁻¹ (wet). All data are averages unless stated.

Site/location	²³⁴ U	²³⁵ U	²³⁸ U
Drigg (Table 4.10)	9.9	0.37	9.5
Ravenglass (Table 4.12)	18	0.60	16
Springfields (Table 4.12)	31	1.5	30
Capenhurst (Table 4.16)	9.5	0.40	9.5
Derby (Table 9.1)	27	0.97	24

90, Section 6.3

The maximum dose due to gaseous disposals was received by adults.

161, Appendix 4

The 1 year old child dose coefficient for ⁹⁹Tc was 4.80 10⁻⁹.

RIFE-4
1998

Page, Section	Comment																
70, Table 4.12	The concentrations of total Cs and ¹⁴⁴ Ce in ovine muscle (max) were 0.61 and <1.8 Bq kg ⁻¹ (wet) respectively. No value for ¹⁵⁵ Eu is available.																
75, Table 4.15(a) 77, Table 4.16 116, Table 9.1	The following activity in soil data were reported as being Bq kg ⁻¹ (dry) whilst they should have been reported as Bq kg ⁻¹ (wet). All data are averages unless stated.																
	<table border="1"> <thead> <tr> <th>Site/location</th> <th>²³⁴U</th> <th>²³⁵U</th> <th>²³⁸U</th> </tr> </thead> <tbody> <tr> <td>Springfields (Table 4.15(a))</td> <td>72</td> <td>3.0</td> <td>68</td> </tr> <tr> <td>Capenhurst (Table 4.16)</td> <td>7.9</td> <td>0.30</td> <td>7.4</td> </tr> <tr> <td>Derby (Table 9.1)</td> <td>31</td> <td>0.93</td> <td>26</td> </tr> </tbody> </table>	Site/location	²³⁴ U	²³⁵ U	²³⁸ U	Springfields (Table 4.15(a))	72	3.0	68	Capenhurst (Table 4.16)	7.9	0.30	7.4	Derby (Table 9.1)	31	0.93	26
Site/location	²³⁴ U	²³⁵ U	²³⁸ U														
Springfields (Table 4.15(a))	72	3.0	68														
Capenhurst (Table 4.16)	7.9	0.30	7.4														
Derby (Table 9.1)	31	0.93	26														
96, Table 6.4(a)	The concentration of ²⁴¹ Am in mud at Paddy's Hole was <1.0 Bq kg ⁻¹ (dry). No measurement of ^{239/240} Pu was made.																
125, Section 11.1	Last but one paragraph. The estimated dose was 0.094 mSv.																
131, Section 11.8	Last paragraph, first sentence. Replace 1997 with 1998.																

RIFE-5
1999

71, Table 4.15(a) 73, Table 4.16 118, Table 9.1	The following activity in soil data were reported as being Bq kg ⁻¹ (dry) whilst they should have been reported as Bq kg ⁻¹ (wet). All data are averages unless stated.																
	<table border="1"> <thead> <tr> <th>Site/location</th> <th>²³⁴U</th> <th>²³⁵U</th> <th>²³⁸U</th> </tr> </thead> <tbody> <tr> <td>Springfields (Table 4.15(a)) max</td> <td>180</td> <td>15</td> <td>200</td> </tr> <tr> <td>Capenhurst (Table 4.16) max</td> <td>12</td> <td>0.46</td> <td>12</td> </tr> <tr> <td>Derby (Table 9.1) max</td> <td>34</td> <td>1.3</td> <td>31</td> </tr> </tbody> </table>	Site/location	²³⁴ U	²³⁵ U	²³⁸ U	Springfields (Table 4.15(a)) max	180	15	200	Capenhurst (Table 4.16) max	12	0.46	12	Derby (Table 9.1) max	34	1.3	31
Site/location	²³⁴ U	²³⁵ U	²³⁸ U														
Springfields (Table 4.15(a)) max	180	15	200														
Capenhurst (Table 4.16) max	12	0.46	12														
Derby (Table 9.1) max	34	1.3	31														
112, Section 8.2	The second sentence of paragraph three states that "the duck and tide washed pasture pathways gave doses of 0.032 and 0.009 mSv y ⁻¹ respectively." The dose due to the duck pathway should read 0.042 mSv y ⁻¹ . The value for tide washed pasture is correct.																
123, Table 10.2	The concentration of ¹⁴ C in grass from Billingham was 960 Bq kg ⁻¹ (wet).																
162, Table A1.2	The Dounreay (Fast Reactor) data were duplicated.																

RIFE-6
2000

31, Section 3.5	It was stated that the dose limits do not apply to natural radionuclides. This sentence should be deleted.												
75, Table 4.16 124, Table 9.1	The following activity in soil data were reported as being Bq kg ⁻¹ (dry) whilst they should have been reported as Bq kg ⁻¹ (wet). All data are averages unless stated.												
	<table border="1"> <thead> <tr> <th>Site/location</th> <th>²³⁴U</th> <th>²³⁵U</th> <th>²³⁸U</th> </tr> </thead> <tbody> <tr> <td>Capenhurst (Table 4.16) max</td> <td>8.5</td> <td>0.35</td> <td>8.4</td> </tr> <tr> <td>Derby (Table 9.1) max</td> <td>24</td> <td>0.96</td> <td>23</td> </tr> </tbody> </table>	Site/location	²³⁴ U	²³⁵ U	²³⁸ U	Capenhurst (Table 4.16) max	8.5	0.35	8.4	Derby (Table 9.1) max	24	0.96	23
Site/location	²³⁴ U	²³⁵ U	²³⁸ U										
Capenhurst (Table 4.16) max	8.5	0.35	8.4										
Derby (Table 9.1) max	24	0.96	23										

Page, Section	Comment
155, Table 12.1	Target date for project 'Tritium and carbon-14 in seafood' should have been March 2003.
166, Table A1.1	Discharges of tritium from Devonport (pipeline) given as 0.87 TBq should have been 0.087 TBq.
168, Table A1.2	Sellafield Discharge limits of alpha and beta activity should have been 0.00196 and 0.328 TBq. Percentage of limit for alpha and beta activity should have been 4.0 and <1. Discharges of tritium and ¹⁴ C from Sellafield given as 213 and 2.58 TBq should have been 355 and 2.94 TBq. Relevant percentages given as 15 and 30 should have been 25 and 34.
RIFE-7 2001 71, Table 4.8 80, Table 4.15(a) 93, Table 5.2(a) 122, Table 7.3 127, Table 8.2(a) 130, Table 9.1	The following activity in soil data were reported as being Bq kg ⁻¹ (dry) whilst they should have been reported as Bq kg ⁻¹ (wet). All data are averages unless stated.

Site/location	⁶⁰ Co	¹⁰⁶ Ru	¹²⁵ Sb	¹³⁴ Cs	¹³⁷ Cs	²³⁴ U	²³⁵ U	²³⁸ U	²⁴¹ Am
Sellafield (Table 4.8)	<0.80	<3.1	<1.1		80				5.8
max	1.2				97	9.3	0.34	9.1	6.0
Springfields (Table 4.15(a))						95	4.6	89	
max									
Harwell (Table 5.2(a))	<0.40			<0.40	2.9				
Featherstone position A (Table 7.3)						9.5	0.41	9.0	
Featherstone position B (Table 7.3)						7.3	0.34	7.5	
Cardiff (Table 8.2(a))				<0.33	5.6				
max				<0.40	6.5				
Derby (Table 9.1)						18	0.80	18	
max						30	1.3	29	

176, Table A1.1 Discharges of Alpha for Hunterston 'A' given as 0.14 TBq should have been 1.4 10⁻⁵ TBq. The % of limit given as 350 should have been <1.

181, Table A1.2 Dungeness 'A' discharge limit and % of limit for tritium should have been 3 and 23 respectively.

RIFE-8
2002
59, Table 4.1 Two tritium results were omitted.
The data are attached.

Table 4.1. Beta/gamma radioactivity in fish from the Irish Sea vicinity and further afield, 2002

Location	Material	No. of sampling observations	³ H
Liverpool Bay	Flounder	2	<25
Mersey estuary	Flounder	2	<25

79, Table 4.14
82 Table 4.17
128, Table 7.1(a)
138, Table 8.2(a)

The following activity in soil data were reported as being Bq kg^{-1} (dry) whilst they should have been reported as Bq kg^{-1} (wet). All data are averages unless stated.

Site/location	^{60}Co	^{106}Ru	^{125}Sb	^{134}Cs	^{137}Cs	^{234}U	^{235}U	^{238}U
Sellafield (Table 4.14)	<0.80	<2.3	<1.2	68				
max	1.0	<2.7	<1.4	82				
Drigg (Table 4.17)								
max						6.9	0.30	6.5
Aldermaston (Table 7.1(a))								
max						8.7	0.35	8.3
Cardiff (Table 8.2(a))				<0.30	6.4			
max					8.1			

102, Figure 6.1

An incorrect bar in Figure 6.1 for Bradwell (2002) is corrected below




Figure 6.1. Caesium-137 concentration in sediments near nuclear power stations

RIFE-1 - RIFE-8
1995-2002

Urenco Capenhurst have reassessed atmospheric discharges of uranium; the reassessed discharges are listed in Table E1.

Table E1. Reassessed atmospheric discharges of uranium from Urenco Capenhurst

Year	Original reported discharge TBq	Reassessed discharge TBq
1993	$1.74 \cdot 10^{-9}$	$2.41 \cdot 10^{-7}$
1994	$6.74 \cdot 10^{-9}$	$2.63 \cdot 10^{-7}$
1995	$2.69 \cdot 10^{-8}$	$2.75 \cdot 10^{-7}$
1996	$1.11 \cdot 10^{-7}$	$8.23 \cdot 10^{-7}$
1997	$6.80 \cdot 10^{-8}$	$4.90 \cdot 10^{-7}$
1998	$6.87 \cdot 10^{-8}$	$1.87 \cdot 10^{-6}$
1999	$8.15 \cdot 10^{-8}$	$1.01 \cdot 10^{-6}$
2000	$9.64 \cdot 10^{-8}$	$8.72 \cdot 10^{-7}$
2001	$1.20 \cdot 10^{-7}$	$9.77 \cdot 10^{-7}$
2002	$1.16 \cdot 10^{-7}$	$6.01 \cdot 10^{-7}$

RIFE-9
200382, Table 3.15
138 Table 6.1(a)
141, Table 6.3(a)
151, Table 7.3(a)
157, Table 8.1(a)The following activity in soil data were reported as being Bq kg⁻¹ (dry) whilst they should have been reported as Bq kg⁻¹ (wet). All data are averages unless stated.

Site/location	⁶⁰ Co	¹⁰⁶ Ru	¹²⁵ Sb	¹³⁴ Cs	¹³⁷ Cs	¹⁵⁴ Eu	²³⁴ U	²³⁵ U	²³⁸ U	²⁴¹ Am
Sellafield (Table 3.15)	<0.90	<3.3	<1.2	<0.40	75	<0.50				5.9
max	1.6	<4.2	<1.6		89	<0.60	11	0.54	10	7.7
Aldermaston (Table 6.1(a))							11	0.48	11	
max										
Derby (Table 6.3(a))							47	1.6	40	
max										
Cardiff (Table 7.3(a))				<0.40	8.8					
max					11					
Drigg (Table 8.1)							6.7	0.26	6.7	
max										

185, Table 9.12

Some data were incorrect. The amended version of the table is attached.

Table 9.12. Concentrations of radionuclides in rainwater and air 2003

Location	Sample	No. of sampling observations	Mean radioactivity concentration ^a in rainwater and air								
			³ H ⁷	Be	⁹⁰ Sr ^b	¹³⁷ Cs	²¹⁰ Pb	²¹⁰ Po	²²⁸ Th	Gross alpha ^b	Gross beta ^b
Ceredigion Aberporth	Rainwater	12	<2.4	<1.6		<0.053	0.10			*	
	Air	4		0.0022		<0.00000052	0.00017			*	
Co. Down Conlig	Rainwater	4		<1.5		<0.022	*			*	
	Air	4		0.0022		<0.00000063	0.00015			*	
Dumfries and Galloway Eskdalemuir	Rainwater	4	4	<2.7	1.2		<0.0098	0.094		*	
	Air	4		0.0018		<0.00000043	0.00013			*	
North Yorkshire Dishforth	Rainwater	4		<2.2		<0.039	*			*	
	Air	4		0.0016		<0.00000055	0.00014			*	
Oxfordshire Chilton	Rainwater	12		<1.5	<0.00064	<0.032	0.32			*	0.074
	Air	13		0.0018		<0.00000034	0.00027	<0.000014		*	0.17
Shetland Lerwick	Rainwater	4		1.6		<0.017	*			*	
	Air	4		0.0015		<0.00000052	0.00010			*	
Suffolk Orfordness	Rainwater	4	<2.2	<2.4		<0.048	*			5.2	
	Air	4		0.0022		<0.00000053	0.00020			*	

* Not detected by the method used

^a Bq l⁻¹ for rainwater and Bq kg⁻¹ for air^b Annual bulk analysis

187, Table 9.14

The concentration of ²¹⁰Po in Cornwall, River Fowey was <0.0098 Bq l⁻¹.

Table 9.16. Estimates of maximum radiation exposure from radionuclides in drinking water, 2003^a

Country	Exposure, mSv Man-made radionuclides ^b	Natural radionuclides ^c	All radionuclides
England	<0.001	0.028	0.028
Northern Ireland	<0.001	0.026	0.026
Scotland	<0.001	^d	^d
Wales	<0.001	0.027	0.027

^a The maximum dose is selected for each nuclide group from data for individual sampling locations.

Many estimates of dose are based on concentration results at limits of detection.

^b Including tritium

^c Including carbon-14

^d Analysis of natural radionuclides was not undertaken

214, Table A1.2

The data shown for Faslane are a duplication of the data for Rosyth and were included in error.

RIFE-10
2004

75, Table 3.7

The entry for Haverigg should read 0.087.

45, Figure 3.8

An incorrect bar in Figure 3.8 for Americium discharge is corrected below:




Figure 3.8. Americium-241 and liquid discharge from Sellafield and concentrations in cod*, lobsters and winkles near Sellafield (* estimated in 2004 due to lack of availability of cod)

87, Table 3.15

151 Table 6.1(a)

154, Table 6.3(a)

166, Table 7.3(a)

173, Table 8.1(a)

The following activity in soil data were reported as being Bq kg⁻¹ (dry) whilst they should have been reported as Bq kg⁻¹ (wet). All data are averages unless stated.

Site/location	⁶⁰ Co	¹⁰⁶ Ru	¹²⁵ Sb	¹³⁴ Cs	¹³⁷ Cs	²³⁴ U	²³⁵ U	²³⁸ U
Sellafield (Table 3.15)	<0.43	<1.4	<0.73					
max	0.80	<1.5	<0.80			16	0.64	15
Aldermaston (Table 6.1(a))								
max						7.8	0.29	7.2
Derby (Table 6.3(a))								
max						27	0.94	23
Cardiff (Table 7.3(a))				<0.47	7.1			
max				<0.50	7.7			
Drigg (Table 8.1)								
max						11	0.42	11

Page, Section	Comment
223, Table A1.1	The % annual limit for ¹⁰⁶ Ru discharge at Sellafield was 7% (not 70%).
246, Table A5.1	Some dose per unit intake values were missing for 1 yr old. These were:

Table A5.1. Dosimetric data

Radionuclide	Dose per unit intake by inhalation using ICRP-60 methodology (Sv Bq ⁻¹)
Sr-90 [†]	1.2E-07
Zr-95 [†]	2.1E-08
Ba-140 [†]	2.6E-08
Pb-210 [†]	4.0E-06
Th-228 [†]	1.4E-04
U-238	9.4E-06

[†] Energy and dose per unit intake data include the effects of radiations of short-lived daughter products

RIFE-11
2005

72, Table 3.3a	Footnote 'd' showed an incorrect value. It should have read: ^d The concentration of ²³⁷ Np was 0.00035 Bq kg ⁻¹
112, Table 4.3a	Column headings should have read: ²³⁹ Pu+ ²⁴⁰ Pu ²⁴¹ Pu
140, Table 5.5a	The result of <0.13 for ²⁴¹ Am in the <i>Fucus vesiculosus</i> samples from Pilot Station was incorrectly put into the ²³⁹ Pu+ ²⁴⁰ Pu column.
206, Figures 9.5 and 9.6	Incorrect units were shown. The correct units were mBq l ⁻¹ .
225, Table 9.15	Incorrect headings in the top part of the table. Should have been as below:

Table 9.15. Concentrations of radionuclides in sources of drinking water in England and Wales, 2005

Location	Sample source	No. of sampling observations	Mean radioactivity concentration, Bq l ⁻¹				
			³ H	⁴⁰ K	⁹⁰ Sr	¹³⁷ Cs	²¹⁰ Po
Wales							
Gwynedd	Cwm Ystradlyn Treatment Works	4	<4.0	<0.020	0.0036	0.0018	<0.010
Mid-Glamorgan	Llwyn-on Reservoir	4	<4.0	<0.045	0.0030	<0.0010	<0.013
Powys	Elan Valley Reservoir	4	<4.0	<0.050	0.0040	0.00090	<0.010

248, Table A1.2	Sellafield discharge limits for alpha and beta should have been 8.90 10 ⁻⁵ and 0.00174 TBq respectively.
251, Table A1.2	Aldermaston Tritium discharge and % limit should have been 14.1 and 8.3 respectively.

RIFE 8-11
2002-2005

Concentrations in sediments	For sediment samples with unusually high water contents it was discovered in 2007 that the resulting sample bulk densities were outside the instrument calibration range. Following investigations a correction factor has been calculated and this has been applied to the affected data from 2002-2005 and the new results are reported here in Table E2.
-----------------------------	---

These amendments do not significantly affect any assessments, charts or statements in the relevant RIFE reports.

Table E2. Amended concentrations of radionuclides in sediment, 2002-2005

Year	Site	Location	No. of sampling observations	Mean radioactivity concentration (dry), Bq kg ⁻¹						
				⁵⁷ Co	⁶⁰ Co	⁶⁵ Zn	⁹⁵ Zr	⁹⁵ Nb	¹⁰⁶ Ru	¹²⁵ Sb
2002	Aldermaston	Reading (Kennet)	4							
		Stream draining south	4							
	Bradwell	Maldon	2		<3.4					
		Waterside	2		<4.0					
	Capenhurst	Rossmore (4.3 km downstream)	2							
	Cardiff	Canal	2							
		West of pipeline	2							
	Devonport	Lopwell	2		<3.7					
	Dungeness	Pilot Sands	2		<0.90					
	Harwell	Appleford	4		<0.60					
Day's Lock		4		<0.50						
Sellafield	Caerhun	2		<3.3		<9.6	<7.7	<23	<9.2	
2003	Aldermaston	Reading (Kennet)	4							
		Aldermaston	4							
	Amersham	Outfall (Grand Union Canal)	3	<0.30	<1.1	<1.5				
	Bradwell	Waterside	2		<2.0					
	Cardiff	Canal	1							
	Derby	River Derwent (downstream)	4		<1.0					
Devonport	Lopwell	2		<2.5						
2004	Aldermaston	Reading (Kennet)	4							
		Aldermaston	4							
		Stream draining south	4							
	Amersham	Upstream of outfall (Grand Union Canal)	2	<6.4	<1.8	<4.1				
	Cardiff	Canal	2		<1.6		<4.5	<2.2	<12	<13
Sellafield	Caerhun	2								
2005	Aldermaston	Reading (Kennet)	4							
	Amersham	Upstream of outfall (Grand Union Canal)	2	<5.3	<1.6	<3.6				
	Cardiff	Canal	2							
		Lydebank Brook	4		<1.7					
	Harwell	Appleford	4		<2.5					
		Appleford	4		<2.5					
	Sellafield	Caerhun	2		<2.6		<8.8	<6.8	<20	<20
Trawsfynydd	Bailey Bridge	2		<8.3					<44	

Year	Site	Location	No. of sampling observations	Mean radioactivity concentration (dry), Bq kg ⁻¹							
				¹²⁵ I	¹³¹ I	¹³⁴ Cs	¹³⁷ Cs	¹⁴⁴ Ce	¹⁵⁴ Eu	¹⁵⁵ Eu	²⁴¹ Am
2002	Aldermaston	Reading (Kennet)	4				7.3				<1.9
		Stream draining south	4				<5.1				<1.2
	Bradwell	Maldon	2			6.5	80				<4.0
		Waterside	2			3.9	59				<13
	Capenhurst	Rossmore (4.3 km downstream)	2				<4.4				
	Cardiff	Canal	2	<0.80			2.4				
		West of pipeline	2	<3.1			33				
	Devonport	Lopwell	2				7.7				
	Dungeness	Pilot Sands	2				<0.90				<1.6
	Harwell	Appleford	4				<13				
Day's Lock		4				6.0					
Sellafield	Caerhun	2			<3.4	430	<25	<7.3	<8.0	75	
2003	Aldermaston	Reading (Kennet)	4				8.0				<1.6
		Aldermaston	4				6.3				<2.7
	Amersham	Outfall (Grand Union Canal)	3	<1.0	<550		<2.1				
	Bradwell	Waterside	2				35				<2.7
	Cardiff	Canal	1	<1.4			16				
	Derby	River Derwent (downstream)	4								
Devonport	Lopwell	2				<10					
2004	Aldermaston	Reading (Kennet)	4				5.4				<1.1
		Aldermaston	4				<3.9				<1.3
		Stream draining south	4				<2.8				1.6
	Amersham	Upstream of outfall (Grand Union Canal)	2	<0.80	<1.4		10				
	Cardiff	Canal	2	<1.5			11				
Sellafield	Caerhun	2			<1.5	220	<5.7	<7.3	<3.1	51	
2005	Aldermaston	Reading (Kennet)	4				<3.9				6.5
	Amersham	Upstream of outfall (Grand Union Canal)	2	<1.0	<9.1		6.2				
	Cardiff	Canal	2	<1.8			9.1				
		Lydebank Brook	4				9.0				
	Harwell	Appleford	4				<11				
		Appleford	4				<11				
	Sellafield	Caerhun	2			<2.5	230	<9.3	<12	<5.3	59
Trawsfynydd	Bailey Bridge	2			<4.2	920				76	

	Page, Section	Comment
RIFE-12 2006	70, Table 2.17	The concentration of ^{241}Am in winkles at Drigg should have been 29.
	103, Section 4 Key points	Line 22 second column replace with • At Dungeness, dose from gaseous discharges increased.
	187, Figure 8.5	The range in the key should have been 2 to 8.
RIFE-13 2007	127, Table 4.5a	The ^{210}Po and ^{210}Pb results are the wrong way round for South Gare winkles. ^{210}Po should be 11 and ^{210}Pb should be 0.46 Bq kg^{-1}
	153, Table 5.1	Derby, the total exposure and exposure from intakes of sediment and water should have been $<0.005 \text{ mSv}$.
	161, Section 6 Key points	Line 17 second column should read • The <i>total dose</i> of 0.008...
	239, Appendix 5	Line 3 first column should read ... indicated that it was likely there would be no adverse impact